


Erste Ergebnisse aus dem Wildschweinmonitoring mittels Drohne und Wärmebildkamera

Dr. Marcel Holy
Natur- und Umweltschutzvereinigung Dümmer e.V.




Wildschäden




Wildschäden




Seuchengefahr



Großvieheinheiten (GVE) je ha LF



Seuchengefahr




MAST | 29.09.2014

agrarheute.com

Afrikanische Schweinepest: Tierärzte in Sorge

Berlin - Tierärzte sind in Sorge. Die Gefahr, dass die Afrikanische Schweinepest Deutschland erreicht, steigt. Die Bundestierärztekammer verabschiedet daher eine Resolution, denn ein Ausbruch hätte verheerende Folgen.



Lebensraum Verlandungszone

Über 200 ha Schilf, Weidengebüsch und Erlenbruch als (Dauer-)Einstand



Lebensraum Verlandungszone



Lebensraum Verlandungszone




Lebensraum Verlandungszone



Bestand/Bestandsentwicklung?

Strecke Hegeringe Damme und Lemförde



Bestand/Bestandsentwicklung?



Bestand/Bestandsentwicklung?




Bestand/Bestandsentwicklung?






















Nutzung der gesamten
Verlandungszone am
Nord-, West- und
Südufer durch
Wildschweine

Ausblick

- Neuregelung der Bejagung des NATURA-2000-Gebiets Dümmer
- Intensivierung der Wildschweinjagd
 - Zur Umsetzung von Schutzverpflichtungen
 - Zur Reduzierung von Konflikten und Gefahren



A photograph of two wild boars (Sus scrofa) lying in a field of tall, dry, yellowish-brown grass. The boars are positioned on either side of the word 'Ende', which is centered in the image. The boar on the left is looking towards the camera, while the one on the right is looking slightly to the right. The grass is dense and reaches up to the boars' heads.

Ende